

RESEARCH PAPER

Development of Value Added Food Products from Tropical Tubers

Parvathi, S.^{1*}, Nithya², Umamaheshwari, S.³ and Subbulakshmi, B.⁴

¹Post-Harvest Technology Centre, TNAU, Coimbatore, Tamil Nadu, India

²&³Dept. of Food Science Nutrition, TNAU, Coimbatore, India

⁴Post-Harvest Technology Centre, TNAU, Coimbatore, Tamil Nadu, India

*Corresponding author: parvathi_in1957@yahoo.com

Paper No. 114

Received: 7-2-2016

Accepted: 5-4-2016

Abstract

A study was conducted to develop value added food products from tropical tubers. Among the minor tubers, *Amorphophallus paeoniifolius*, *Typhonium trilobatum* and *Colocasia esculenta* were selected and subjected to different pre-treatments and standardized value added products. The sensory evaluation and nutritional analysis showed that the products developed by using tubers were acceptable up to 30 per cent incorporation. The standardized products from the selected tubers can be explored cottage industries for income generation and employment opportunities. Tuber crops have a prominent role in feeding the world and these crops are going to be an important component in food security system in the coming decades.

Keywords: *Amorphophallus paeoniifolius*, *Typhonium trilobatum*, *Colocasia esculenta*, Tropical tubers, value added

Roots and tubers are the third important food for human after cereals grain and legumes. These crops constitute either staple or subsidiary food for about one fifth of the world population and are known to supply a cheap source of energy especially for the weaker sections of the population. India's population having touched 100 million, still depends on agriculture as the primary occupation. In spite of the increasing production and availability of food grains, there will be a net shortage of nearly 26 million tones by 2020 which is equivalent of 80 tones of roots and tubers (Edison, 2006).

The ever increasing population level coupled with rapidly shrinking cultivable area and increasing fragile resources lead to inadequate supply of food to the people. In order to meet this unfortunate scenario, tuber crops constitute the important link

to fit the food security gap and alleviate hunger and combat malnutrition including nutritional disorders to a sizeable percentage. The nutritional disorders are due to insufficient intake of protein, vitamin A, vitamin C and calcium which could be easily alleviated by adequate supply and consumption of roots and tubers (Padmaja *et al.*, 2013).

The importance and dietary fibre in tubers is well recognized. Dietary fibre reduces the serum cholesterol, prevents colon cancer and maintains good intestinal health as well as provides prophylactic action for cardio-vascular diseases, diabetes and obesity (Pugalendhi *et al.*, 2013). The tuber crops were broadly classified as temperate and tropical groups. Potato is primarily a temperate crop, whereas the tropical tuber crops are tapioca (cassava), sweet

potato, aroids (elephant footyam, taro, tannia and gaint taro), yams (greater yam, lesser yam, african yam) and other minor tubers namely chinease potato, arrow root and yam bean.

Among the aroids, elephant footyam (*Amorphophallus paeoniifolius*) Bengal arum *Karunai kilangu* (*Typhonium trilobatum*) and taro or old cocoyam (*Colocasia esculenta*) are an important tropical minor tuber crops. As a primary staple, they form the major energy source for a vast segment of population in South Pacific and Africa. As a secondary staple or vegetable, these crops find wider acceptability in most of the countries including India. Nutritionally, these tubers are rich in carbohydrate, vitamins and minerals.

But the minor tubers under aroids such as elephant footyam, Bengal arum (*Karunai kilangu*) and taro are not exploited for industrial uses, are cultivated in small pockets and used only as vegetables. To make them important component in our food security systems of this minor tubers, it is necessary to develop value added products. Development of value added products of these tubers to enhance their potential uses within the food industry for the replacement of traditional forms and carbohydrates, and produce entirely new products from the selected tubers such as elephant footyam, *Karunai kilangu* and taro.

Materials and Methods

Collection of selected tubers

Good quality tubers of elephant footyam, bengal arum-karunai *kilangu* (*Typhonium trilobatum*) and taro (*Colocasia esculenta*) were purchased from local market from Madurai district, Tamil Nadu, India for processing into flour and also for standardization of value added products.

Pre-processing of selected tubers for prevention of browning and aridity

The selected tubers were cleaned well under running water for three to four times to remove the adhering sand on the skin and damaged parts

if any. Af er washing, the tubers were surface dried and peeled by using knife.

In the selected tubers, browning and acidity are the major problem. For prevention of browning and aridity from the selected tubers, different treatments were tried namely blanching for 5 min (T_1), soaking at 1 per cent potassium metabisulphate for 30 min (T_2), soaking in 1 per cent citric acid for 30 min (T_3), soaking in 2 per cent hydrochloric acid for 30 min (T_4), soaking in 2 per cent tartaric acid for 30 min (T_5), soaking in 2 per cent citric acid for 30 min (T_6), soaking in 2 per cent salt in cold water (T_{7a}), and hot water for 30 mins (T_{7b}), soaking in 2 % baking soda in cold water (T_{8a}), and hot water for 30 min (T_{8b}), soaking in 2 per cent tamarind juice (3 g in 5 ml of water) for 30 min (T_9), soaking in 2 per cent lime juice for 30 min (T_{10}), steaming for 10 min (T_{11}), open cooking (500 ml water for 10 min) (T_{12}), pressure cooking 5 min (T_{13}), soaking in but er milk (300ml) over night (T_{14}), soaking in 1 per cent citric acid for one hour (T_{15}) and soaking in 1 per cent citric acid for two hours (T_{16}).

Processing of selected tubers into flour

Af er the pre-treatment, the tubers were washed in tap water and cut into (1-2 mm) cubes. The cubes were blanched for 5 min, blanching helps to kill the harmful microorganisms, sof en the cell walls, speed up the drying process and increase the shelf life and to prevent fresh odour in the processed tuber flour.

Af er blanching, the tuber cubes were spread on the trays and were dried in the dehydrator. The cabinet drier was pre-heated at 80° C and the trays were loaded with tuber cubes which was kept in the cabinet dryer. The temperature was maintained at 80° C for 4 hours. The dried cubes were subjected to milling process by using pulverizer, to obtain the fine flow that was sieved by using the sieve (SSS. No. 80). The details of preparation of flour from the selected tubers are given in figure 1.




Fig. 1: Processing of tuber in the form of flour

Standardization of value added products from the selected tubers

(a) Puffed snack

Preparation of puffed snack was standardized by incorporating (elephant foot yam, *karunai kizhangu* and Taro) flour at different level of 10, 20 and 30 percent. The details of processing of puffed snack are given in Table 1.

Table 1: Standardization of puffed snack

Ingredients	C*	T1	T2	T3
Tuber flour (g)	-	10	20	30
Maize flour (g)	100	90	80	70
Pepper powder (g)	2	2	2	2
Water (ml)	10	10	10	10
Salt	2	2	2	2

C* = Control




Fig. 2: Processing of puffed snack

(b) Papad

Papad was standardized using (elephant foot yam, *karunai kizhangu* and Taro) flour at different incorporation of 10, 20, and 30 per cent. The details of processing of papad is given in Table 2.

Table 2: Standardization of papad

Ingredients	C*	T1	T2	T3
Tuber flour (g)	—	10	20	30
Black gram flour (g)	100	90	80	70
Salt	5	5	5	5
Water (ml)	40	40	40	40
Sodium bicarbonate (g)	1	1	1	1
Oil (g)	5	5	5	5

C* = Control

Preparation of papad




Fig. 3: Preparation of papad

Sensory evaluation of standardized value added products

To ensure the acceptability of the modified recipes used for products, they were subjected to evaluation by composite scoring for their sensory qualities. The standardized value added products were prepared and presented to a panel of 15 judges. Specific sensory characteristics of each product (appearance, color, flavor, taste, texture and overall acceptability) were rated separately using hedonic scale on a scale of 1 to 9. Scores were defined as follows: 1 – dislike extremely, bad; 9 – like extremely, excellent. Numerical averages were then, calculated for a composite test score (Joshi, 2006).

Physical properties of standardized value added products

(a) Puffed snack

Bulk Density

The bulk density was calculated by measuring the actual dimensions of extrudates (AACC, 2000). The diameter and length of extrudates were measured using Vernier caliper. The weight per unit length of extrudate was determined by weighing a measured length (about 1 cm). The bulk density was then, calculated using the following formula, assuming a cylindrical shape of extrudate.

$$\rho b = \frac{4}{\pi l d^2}$$

Where *b* is bulk density (g/cm³), *l* is the length per gram of the extrudates (cm/g) and *d* is diameter of the extrudates (cm). Five pieces of extrudates were randomly selected and average taken.

Water solubility index and water absorption index

Water solubility (WSI) and water absorption indices of extruded products were determined by a modification of the method of Anderson *et al.* (1969). The extrudates samples were ground and sieved through 500 sieves. Distilled water (10 ml) at 25° C was placed in a tarred centrifuge tube and 0.5 g of extrudate was dispersed

in the water. Care was taken to avoid lumping in order to produce smooth dispersion. After standing for 30 min (with intermittent shaking every 5 min), the sample was centrifuged (Roto Silenta II, Hettich, Tuttingen, Germany) at 1800 × g for 15 min. The supernatant was decanted into a tarred aluminum pan and dried at 105°C until got a constant weight. The weight of the gel remaining in the centrifuge tube was noted and the results are expressed as the average of two measurements.

(b) Papad

The weight, diameter and thickness of the papad were determined. Weight of the papad was measured using an electronic balance (model: OPAL ISO 9001: 2008 certified) having least of 10 mg. Diameter was assessed having a least count of 0.01 mm thickness which was calculated by using screw gauge having count of 0.1 mm.

Nutrient analysis of value added products from minor tubers

Based on the acceptability score, the standardized value added products from minor tubers were analysed for its proximate composition using AOAC (1984) method.

Estimation of calcium oxalate

Edible aroid (*Colocasia*, and *Amorphophalles*) accumulate oxalic acid in the form of free oxalate and bound oxalate (mainly as calcium oxalate). The presence of calcium oxalate has been attributed to be one of the major causes of acidity. Hence to prevent acidity, different pre-treatments were tried as detailed earlier. Among the 16 treatments tried, soaking in citric acid at 1 per cent level for one hour was found to be effective in preventing browning and acidity in elephant foot yam and taro flour by organoleptic evaluation. For Bengal arum (*karunai kizhangu*), The tubers treated with citric acid at 1 per cent for 2 hours was found to be effective in preventing the acidity in the preliminary results. Hence the tubers treated with citric acid were used for standardizing different value added products.

The presence of calcium oxalate was estimated for the raw (unprocessed) and pretreated flour with citric acid as per the procedure suggested in the AOAC method (1984).

Results and Discussion

Pre-processing of selected tubers

The selected tubers soaked in 1 per cent citric acid for one hour (T_{15}) were effective in elephant foot yam and taro. For Bengal arum (*karunai kizhangu*), the tubers soaked in 1 per cent citric acid for 2 hours (T_{16}) was found to be effective in reducing the acidity and browning. It was found that processed flour had bright color and good aroma in organoleptic evaluation. Sundaresan (2005) reported that citric acid and tartaric acid at 1 per cent concentration were found to be effective in reducing the acidity and oxalate levels. However, citric acid was found to be a better choice as it imparted a bright colour to the cooked sample. Soaking of tubers in solvents prior to boiling was also found to be equally effective in reducing the acidity. Nearly 75% calcium oxalate was lost in treatment with citric acid and tartaric acid.

Estimation of calcium oxalate content in fresh and processed tuber flour

The calcium content in the fresh and pre-processing tuber flour was analyzed. The details are presented in table 3.

Table 3: Presence of Calcium oxalate content in fresh and processed tuber flour

Tuber	Calcium oxalate	
	Fresh tuber flour	Pre-processed tuber flour
Elephant foot yam	0.14	0.05
Bengal arum (<i>karunai kizhangu</i>)	0.25	0.09
Taro	0.20	0.08

From the results it was found that the calcium oxalate content in fresh tuber were 0.14% in elephant foot yam, in *karunai kizhangu* 0.25% and in taro 0.20

per cent, respectively. The processed (soaking the tubers in citric acid followed by blanching, drying & roasting) tuber flour was subjected for the presence of calcium oxalate. It was found that elephant foot yam flour had 0.05%, *karunai kizhangu* flour contained 0.09% and in taro flour had 0.08 percent calcium oxalate. The reduction in acidity is clearly evident from the results.

The reduction in oxalate content may be due to different treatment like peeling, washing, dicing, soaking, blanching and roasting. Akapan and Umoh (2004) stated in their study that the peel of tubers contain more oxalate than the peeled tubers. Buntha *et al.* (2008) also reported that the acidity of taro can be reduced by peeling, grating, soaking and fermentation. Huang *et al.* (2007) also found that soaking reduced the concentration of oxalates by 23.5% and by cooking 56.7% Shanthai kumari *et al.* (2008) further confirmed that the decrease in the anti-nutritional factors like oxalates during soaking may be due to leaching out of these substances in soaking medium. Boiling may cause considerable cell rupture and facilitate leakage of soluble oxalate into cooking water as per the results reported by Albihn & Savage (2001).

Iwuo and Kalu (1994) reported that boiling of taro had reduced the oxalate content by 65.7-82.1%, the study conducted by Adane *et al.* (2013) also reported that boiling of taro had reduced the oxalate content by 70.9%. Sundaresan (2005) found that in fresh tuber of taro cultivars 0.07-0.15% was calcium oxalate content, elephant foot yam had 0.12-0.24 per cent. It is concluded that addition of mild organic acid prior to boiling was found to be effective in reducing the acidity more than 75 per cent.

Incorporation of organic acid in cooking medium provides low pH and facilitate solubilization of calcium oxalate. In the analysis, it was found, that the selected processed tuber flours had 60-64 per cent reduction in calcium oxalate by treating the tubers in citric acid.

Organoleptic evaluation of value added products from minor tubers

(a) Puffed snack

The prepared puffed snack was organoleptically evaluated, to assess the acceptability, the prepared puffed snack was given to 15 trained judges for organoleptic evaluation using nine point hedonic scale. The results of sensory evaluation are given in Table 4 and figure 4.



Fig. 4: Popped snacks

Table 4: Organoleptic evaluation of puffed snack

Particulars	C*(%)	T ₁ (%)	T ₂ (%)	T ₃ (%)
Colour	85	82	80	85
Texture	84	82	75	83
Flavour	85	80	76	85
Taste	85	83	75	85
Overall acceptability	85	83	80	85

C* Control

Based on the sensory evaluation thirty per cent (T₃) incorporation of elephant food yam flour, *karunai kizhangu* flour and taro flour incorporated puffed snack had higher acceptability. The product had the overall acceptability score of 85 per cent. Kavya and Reddy (2011) made an attempt to formulate extruded snacks in combination with corn grits, black gram dhal, root and tuber like potato, sweet potato, colocasia and beet root, corn grits, black

and gram dhal and tubers were added in the ratio of 60: 20: 20 respectively to produce puffed snacks. It was concluded that the most acceptable based extruded products can be prepared by using tubers. The products need to be explored further for commercialization as an innovative snack food to increase the utility of the tubers.

(b) Organoleptic evaluation of papad

The prepared papad was fried in oil and it was organoleptically evaluated, to assess the consumer acceptability of the product. The results of organoleptic evaluation are given in Table 5 and the product is shown in figure 5.



Fig. 5: Papad

Table 5: Organoleptic evaluation of papad

Particulars	C*(%)	T ₁ (%)	T ₂ (%)	T ₃ (%)
Colour	95	84	86	93
Texture	94	84	84	93
Flavour	92	80	80	93
Taste	95	80	74	95
Overall acceptability	95	80	80	90

C* Control

Based on the organoleptic evaluation, thirty percent (T₃) incorporation of elephant foot yam flour, *karunai kizhangu* flour and taro flour incorporated papad had higher acceptability. Papad at 50% incorporation of taro was highly accepted by the judges. But while kneading and spreading, the dough texture was

Table 6: Nutrient content of the standardized tubers products in 100gms

Name of the product	Moisture (%)	Carbohydrate (g)	Protein (g)	Fat (g)	Fiber (g)	β-carotene (μg)
Elephant foot yam						
Puffed snack	23.61	5.52	0.36	0.03	0.24	78
Papad	6.20	45.42	4.90	4.23	3.5	142.1
Karunai kizhangu						
Puffed snack	20.97	7.80	0.42	0.03	0.30	23.4
Papad	6.50	52.10	5.10	3.7	4.10	87.5
Taro						
Puffed snack	21.93	6.33	0.90	0.03	0.30	12.4
Papad	6.40	43.50	3.70	3.50	3.80	64.10

sticky, it may be due to the mucilage content in taro flour. The product had the overall acceptability score of 85 per cent.

Nutrient content of the value added products from the selected tubers

Based on the organoleptic evaluation T₃ sample had higher acceptability score, when compared to other incorporation level of selected tuber flour. Hence proximate composition for the 30 per cent incorporated tuber flour value added products were estimated and presented in Table 6.

The standardized puffed snack was prepared from three different minor tubers such as elephant foot yam, *karunai kizhangu* and taro. The results indicated that moisture content of elephant foot yam product was high compared to *karunai kizhangu* and taro

products. *Karunai kizhangu* flour exhibited maximum carbohydrate content at (7.77 g) followed by elephant foot yam and taro. The values of fat in all the products were equal. Fibre content of *Karunai kizhangu* and taro was equal but was lower in elephant foot yam. Highest amount of beta carotene was obtained from elephant foot yam when compared to *karunai kizhangu* and taro.

The mineral content of the standardized products

From the table it was observed that calcium content was higher in elephant foot yam puffed snack followed by *karunai kizhangu* and taro. Higher amount of phosphorous, magnesium and potassium content were observed in taro products and lower in elephant foot yam. *Karunai kizhangu* had high amount of iron and zinc.

The mineral content of the standardized products are presented in Table 7.

Name of the product	Calcium (mg)	Phosphorus (mg)	Iron (mg)	Magnesium (mg)	Potassium (mg)	Zinc (mg)
Elephant foot yam						
Puffed snack	30.0	251.56	2.01	100.50	197.98	2.3
Papad	119.8	267.7	2.54	88.0	50.24	1.10
Karunai kizhangu						
Puffed snack	23.92	255.60	2.11	104.82	270.0	2.47
Papad	115.3	265.5	2.72	93.1	119.04	1.23
Taro						
Puffed snack	25.7	282.65	1.92	107.98	361.63	2.41
Papad	115.0	295.5	2.5	95.4	210.5	1.1

Papad prepared from elephant foot yam flour contain high amount of moisture followed by *karunai kizhangu* had taro. Carbohydrate content of *karunai kizhangu* papad was high to elephant foot yam and taro. Protein content was more in taro, *karunai kizhangu* and elephant foot yam papads. But there is no change in fat content of papad prepared from the three tubers flour i.e. all the three tubers showed the equal fat content (0.03g). equal amount of fibre content was present in *karunai kizhangu* and taro papads, but in elephant foot yam papads contained less amount of fibre. Beta carotene content of elephant foot yam papad was higher compared to *karunai kizhangu* and taro. Calcium content was higher in elephant foot yam papad. The highest amount of phosphorus, magnesium, potassium were found in taro papads followed by *karunai kizhangu* and elephant foot yam. *Karunai kizhangu* papad contain higher amount of iron and zinc compared to others.

Conclusion

Diversification for the production of value added products is one of the methods to retain the minor tubers in the existing cropping systems. This will directly or indirectly lead to the generation of employment opportunities. The standardized foods from the selected tubers can be exploited at the cottage industries level. It can be concluded that by soaking the selected tubers in one per cent citric acid for one hour for elephant foot yam and 2 hours for taro and *karunai kizhangu* helps to prevent browning and acidity. The processed tubers can be converted into grits, *suji* and flour and it can be used for developing different types of value added products. Further, the selected tubers had the potential to develop puffed snack and papad with 30 percent of tuber flour incorporation.

References

Adane, T., Ashimelis, R., Negussie, B., Tilahunand, G.D. and Haki 2013. Effect of processing method on the proximate composition, minerals content of anti-nutritional factors of taro grown in Ethiopia. *African J Food, Agri. Nutr and Develop.* **13**(2): 73-91.

- Akpan, E.J. and Umoh, IB. 2004. Effect of Heat and Tetracyclic Treatment on the food quality of acidity factors in cocoyam (*Xanthosoma sagittifolium* L Scholt). *Pak.J.Nutr.* **3**: 240-243.
- Albihn, P.B. and Savage, G.P. 2001. The bio availability of oxalate from oca(*oxalis tuberose*) *J. Urol.* **166**: 420-422.
- American Association of Cereal Chemist. 2000 approved methods of the AACC 10th edn.
- Andesson, R.A., Conway H.F., Pferifer V.F and Griffin F. 1969. Gelatinisation of corn grits by roll and extrusion cooking. *Cereal Science Today.* **14**: 4-7.
- AOAC official methods of Analysis. 1984. 14th edition, AOAC Washington DC/ 611-612.
- Buntha, P., Borin, K. Preston, T. and oligle, B. 2008. Survey of taro varieties and other use in selected areas of cambodiya *Livestock Research for Rural Development.* **20**.
- Edison, S. 2006. Tropical tuber crops: Issues & strategies for sustainable development. Souvenir, 14th Triennial Symposium of the international society for Tropical Root Crops. 61-63.
- Huang, C.C., Chen, W.C. and Wang, C.C.R. 2007. Comparison of Taiwan paddy and upland cultivation of taro (*Colocasia esculanta* L) cultivars for nutritive values. *Food Chem.* **102**: 250-256.
- Joshi, V.K. 2006. Sensory science: Principles and Application in Food Evaluation. Agro-tech Publishing Academy, Udaipur (India), p. 527+illustrations.
- Iwuoha, I. and Kalu, A. 1994. Calcium oxalate and physico chemical properties of cocoyam tuber flour as affected by processing. *Food chem.* **54**: 61-64.
- Kavya and Reddy, M. 2011. Development of evaluation of ready to eat extruded snacks by incorporating roots & tubers 43rd National Conference of Nutrition society of India. Organized by National Institute of Nutrition & ICAR in Nov. 11-12.
- Padmaja, G., Sheriff, J.T. and Sajiev, M.S. 2013. Healthy food from tubers. *Indian Horticulture.* **58**(3): May – June: 18
- Pugalendhi, L.M. Prabhu, Venkatesan, K. and Anbukkarasi, V. 2013. Bio diversity of Aroids, Aroids opportunities and challenges CTCRI, Sreekariyam, Thiruvananthapuram Kerala India, II.
- Shanthakumar, S., Mohan, V. and De Brultov. 2008. Nutritional and anti nutritional evaluation of wild yam (*Dioscorea* spp). *Trop Sub Trop Agro Ecosystem* **8**: 313-319.
- Sundaresan. 2005. Acridity of edible aroids, Technical bulletin 38 CTCRI, KAR, T. No 20-21.