

Effect of diluent supplementation with garlic extract on semen quality of cocks during liquid storage

Hazim J. Al – Daraji

Department of Animal Production, College of Agriculture, University of Baghdad
Baghdad

Abstract

This experiment was designed to examine whether garlic extract (GE) supplementation could improve rooster sperm motility, viability, and morphology during *in vitro* storage for different periods (24, 48 or 72 h). A total of 42 White Leghorn roosters, 22 – wk old randomly divided into 6 experimental pens (7 roosters each) were used in this study. The experimental groups were as follows: T1 = fresh , undiluted semen (control); T2 = semen diluted 1:1 with Lake diluent (LD) alone; T3 = semen diluted 1:1 with GE alone, while T4, T5 and T6 represented semen samples diluted 1:1 with LD and supplemented with 1, 2 and 4 ml GE / 100 ml of diluent, respectively. Results denoted that semen incubation for 24, 48 or 72 h at the refrigerator temperature in the absence of GE (T1) was associated with a significant ($p<0.05$) decrease in the mass activity and individual motility, and significant ($p<0.05$) increase in the percentages of dead spermatozoa, abnormal spermatozoa and acrosomal abnormalities. However, the inclusion of GE into the LD (T4, T5 and T6) significantly ($p<0.05$) improved motility, viability and normality of spermatozoa acrosomes compared with control group (T1). Besides , T5 and T6 surpasses all other treatments in ameliorate the deterioration that found in the semen traits included in this study after *in vitro* storage for up to 72 h . In addition, T2 was superior to T3 as regards mass

activity and individual motility, whereas there were no significant differences between these two treatments with relation to percentages of live spermatozoa and normal spermatozoa and acrosomes for semen samples stored for 24, 48 or 72 h. In conclusion, supplementation of GE into avian semen diluents particularly at the doses of 2 and 4 ml GE / 100 ml of diluent can be used as successful technique for depresses the detrimental effects of lipid peroxidation which could lead to sperms deterioration during *in vitro* storage for up to 72 h.

Keywords: Garlic extract, Semen quality, Liquid storage, Cocks

Introduction

Spermatozoa are unique in structure and chemical composition and are characterized by high proportions of polyunsaturated fatty acids (PUFAs) in the phospholipid fraction of their membranes. This characteristic composition confers to sperm plasma membrane the fluidity they require to undergo the membrane fusion events that characterize fertilization (30). However, high level of PUFAs increases the susceptibility of cells to free radical

*Corresponding Author: prof.hazimaldaraji@yahoo.com

MS Received: 1st January, 2013

MS Accepted: 10th March, 2013

attack and lipid peroxidation. Therefore, antioxidant protection is a vital element in maintaining motility, viability, membrane integrity, and fertilizing ability (Al-Daraji, 2004). The major fatty acids components of avian spermatozoa are arachidonic (20: 4 $n - 6$) and docosatetraenoic (22: 4 $n - 6$) acids (Kelso *et al.*, 1996). Thus avian spermatozoa are characterized by high amounts of C₂₀₋₂₂ polyunsaturates of the $n - 6$ series, whereas long - chain fatty acids of the $n - 3$ series predominate in mammalian spermatozoa. However, it appears that the 20: 4 $n - 6$ and 22: 4 $n - 6$ present in avian spermatozoa performs an essential function in promoting optimal spermatozoa motility, viability and fertilizing capacity, as marked reductions in the amounts of these fatty acids in spermatozoa as a result of lipid peroxidation are associated with impaired sperm number, motility, viability and fertilizing ability (Kelso *et al.*, 1997).

Free radicals – atoms formed when oxygen interacts with certain molecules during normal bodily processes or from exposure to several environments. Garlic is one antioxidant that defends against free radicals damage, thereby preserving the body's healthy functioning (Munday *et al.*, 1999). However, garlic antioxidants help scavenges free radicals – particles that can damage cell membranes, interact with genetic material and possibly contribute to the aging process as well as the

development of a number of conditions including heart disease and cancer. Garlic antioxidants can neutralize free radicals and may reduce or even help prevent some of the damage they cause over time. In the study of Silagy and Haw (1994), a total of 261 patents from 30 general practices were given either garlic powder or placebo. After a 12 week treatment period mean serum cholesterol levels dropped by 12% in the garlic treated group and triglycerides and total lipid levels decreased by 17% and 19%, respectively compared to the placebo group .

Because garlic (*Allium sativum* L) has antioxidant activity, this study examines the potential role of garlic extract (GE) as an antioxidant in preserving rooster's spermatozoa during *in vitro* storage for certain storage periods (24, 48 or 72 h).

Material and methods

Forty – two White Leghorn roosters, 22 – wk old randomly divided into 6 experimental pens (7 each) were used in the experiment. Cocks fed a commercial layer ration *ad libitum*. Semen samples were collected from all roosters once a week for 10 consecutive weeks (22 – 32 weeks of age) by using the method of Lake and Stewart (Lake and Stewart, 1978). Semen samples in each treatment pen were divided into 3 test tubes of 1 ml each to provide 3 replicates pooled samples per each treatment group. Therefore, there were 30 replicates for each treatment. The

experimental groups were as follows : T1 = fresh , undiluted semen (control group); T2 = semen diluted 1:1 with LD alone ; T3 = semen diluted 1:1 with GE alone ; T4 = semen diluted with LD and supplemented with GE (1 ml / 100 ml of diluent); T5 = semen diluted with LD and supplemented with GE (2 ml / 100 ml of diluent) and T6 = semen diluted with LD and supplemented with GE (4 ml / 100 ml of diluent). Experimental samples were stored at the refrigerator temperature (4 – 6 °C) for different storage times (24, 48 or 72 h.). An aliquot of semen from each group was evaluated at 24, 48 and 72 h of *in vitro* storage for mass activity, individual motility, and percentages of dead spermatozoa, abnormal spermatozoa and acrosomal abnormalities.


Spermatozoa motility (movement in a forward) was estimated on a percentage basis by using the microscopic method of Sexton (1976). The determination of percentage of dead spermatozoa was done by using a Fast green – stain – Eosin B stain – glutamate extender (Al-Daraji *et al.*, 2002). Percentage of abnormal spermatozoa was evaluated by using a Gentian violet – eosin stain (Al-Daraji, 1998). As an alternative to evaluate the percentage of acrosomal abnormalities, staining procedure for fixed samples have been developed to distinguish which spermatozoa have retained or lost the acrosome (Al-Daraji, 2001). However, the extraction of garlic components was achieved according to the procedure that mentioned by Ohnishi and Ohnishi

(Ohnishi and Ohnishi.2001). Results were evaluated by analysis of variance. Differences between experimental groups' means were analyzed by Duncan's multiple range test, using the ANOVA procedure in Statistical Analysis System (SAS.1996).

Results and discussion


The overall means of the treated groups regarding mass activity and individual motility at 0 h shows that the differences between treated groups and the control group were not statistically significant (Figures 1 and 2). However, in semen samples evaluated after 24, 48 or 72 h *in vitro* storage at the refrigerator temperature, diluent supplemented with GE (T4, T5 and T6) surpasses T1 and T3 groups as regards mass activity and individual motility. However, there were no significant differences between T2 and T4 groups with relation to these two traits when semen samples stored for 24 or 48 h, while T4 was superior to T2 group when samples evaluated after 72 h of storage. Besides, T5 and T6 surpass all other treatments in regard to mass activity and individual motility during all storage periods (Figures 1 and 2).

As shown in Figures 3, 4 and 5, there were no significant differences between T1, T2, T3 and T4 groups, and between T5 and T6 groups as regards percentages of dead and abnormal spermatozoa and acrosomal abnormalities for semen samples evaluated directly after collection (0 h).



T1 = Fresh semen , T2 = LD , T3 = G , T4 = LD + G (1ml / 100 ml) , T5 = LD + G (2 ml / 100 ml) and T6 = LD + G (4 ml / 100 ml) . Bars with different superscript differ significantly (p < 0.05) .

Figure 1: Effect of supplementation of different levels of garlic extract into LD on mass activity of rosters semen stored for different periods.




T1 = Fresh semen , T2 = LD , T3 = G , T4 = LD + G (1ml / 100 ml) , T5 = LD + G (2 ml / 100 ml) and T6 = LD + G (4 ml / 100 ml) . Bars with different superscript differ significantly (p < 0.05) .

Figure 2: Effect of supplementation of different levels of garlic extract into LD on individual motility of rosters semen stored for different periods

However, during this time of storage (0 h) T5 and T6 surpasses other treatments in relation to these three characteristics. In addition, supplementation of the diluent with GE (T4, T5 and T6) significantly improved the percentages of live spermatozoa and normal spermatozoa and acrosomes when semen samples *in vitro* stored for 24, 48 or 72 h. The most efficient levels were 2 ml GE / 100 ml of diluent (T5) and 4 ml GE / 100 ml of diluent (T6). On the other hand, there were no significant differences between T2 and T3 groups regarding these 3 characters (Figures 3, 4 and 5).


It is interestingly obvious from the results of this study that inclusion of GE into LD closely maintained activity, viability and normality of sperms and acrosomes. Our results are in

accordance with the results of previous authors (Al-Daraji, 2000; Al-Daraji, 2002; Al-Daraji, 2004; El-Nasry *et al.*, 2004) who found that the addition of certain antioxidants (vitamins A, C or E) to the avian semen diluents had preserved motility, viability, morphology and fertilizing capacity of semen stored for different storage periods at 4-6°C . Furthermore, the amelioration in semen characteristics noticed in the present experiment may was a result of GE antioxidants repressing or limiting the detrimental effects of lipid peroxidation during *in vitro* storage. Wishart (1989) reported that lipid peroxidation is already initiated in fresh ejaculates and is able to develop during incubation even at low temperatures. Donoghue and Donoghue (1997) pointed out that antioxidant activity in seminal plasma and sperm is




T1 = Fresh semen , T2 = LD , T3 = G , T4 = LD + G (1 ml / 100 ml) , T5 = LD + G (2 ml / 100 ml) and T6 = LD + G (4ml / 100 ml) .
 Bars with different superscripts differ significantly (p < 0.05)

Figure 3: Effect of supplementation of different levels of garlic extract into LD on dead spermatozoa of roosters semen stored for different periods.



T1 = Fresh semen , T2 = LD , T3 = G , T4 = LD + G (1 ml / 100 ml) , T5 = LD + G (2 ml / 100 ml) and T6 = LD + G (4 ml / 100 ml) .
 Bars with different superscripts differ significantly ($p < 0.05$) .

Figure 4: Effect of supplementation of different levels of garlic extract into LD on abnormal spermatozoa of rosters semen stored for different periods .



T1 = Fresh semen , T2 = LD , T3 = G , T4 = LD + G (1 ml / 100 ml) , T5 = LD + G (2 ml / 100 ml) and T6 = LD + G (4 ml / 100 ml) .
 Bars with different superscripts differ significantly ($p < 0.05$) .

Figure 5: Effect of supplementation of different levels of garlic extract into LD on acrosomal abnormalities of rosters semen stored for different periods .

not high enough to prevent lipid peroxide damage after extension and *in vitro* storage, and that supplemental antioxidants could improve semen shelf life.

Extracts of fresh garlic contain antioxidant phytochemicals that prevent oxidant damage. These include unique water-soluble organosulfur compounds, lipid-soluble organosulfur components and flavonoids, notably allixin and selenium (Ohnishi and Ohnishi.2001). A mounting body of research indicates that garlic act as a potent antioxidant, decreasing lipid peroxidation, increasing free radical scavenging and glutathione, lowering high cholesterol by interfering with its metabolism in the liver and lowering LDL (bad) cholesterol and triglyceride levels while raising the level of HDL (good) cholesterol (Geng and Lau.1997; Reeve *et al.*, 1993; Wei and Lau.1998). Ohnishi and Kojima (1997) concluded that aged garlic extract has a strong antioxidant effect. However, GE exerts antioxidant action by scavenging reactive oxygen species (ROS), enhancing the cellular antioxidant enzymes superoxide dismutase, catalase and glutathione peroxidase, and increasing glutathione in the cells (Ryu *et al.*, 2001). GE inhibits lipid peroxidation, reducing ischemic / reperfusion damage and inhibiting oxidative modification of LDL, thus protecting endothelial cells from the injury by the oxidized molecules (Borek, 2001). However, phytochemicals from plant – rich diets, including garlic,

provide important additional protection against oxidant damage (Borek, 1997). The variety of antioxidant phytochemicals in GE, which protect against detriments-causing oxidative damage (Ide and Lau.1997), may act in single and combined fashion (Amagase *et al.*, 1996). The antioxidative actions of GE and its components are determined by their ability to scavenge ROS and inhibit the formation of lipid peroxides. These effects are determined by measuring the decrease in ROS-induced chemiluminescence, inhibition of thiobarbituric acid reactive substances (lipid peroxides), and *in vitro* inhibition of the release of pentane, a product of oxidized lipids, in the breath of an animal exposed to oxidative stress (Imai *et al.*, 1994; Awazu and Horie. 1997). On the other hand, Pizzorno *et al.* (1999) reported that garlic has always been known as an aphrodisiac and from a medical point of view it can improve blood circulation significantly. Now appear that an enzyme called nitric oxide synthase is primarily responsible for the mechanism of erection. Studies have recently shown that garlic in certain forms can stimulate the production of nitric oxide synthase particularly in individuals who have low levels of this enzyme. Clearly folklore is now being proven correct.

In the light of the results of the present study, it can be concluded that developing a defence system against lipid peroxide damages of practical importance to improve the extended liquid storage of roosters' semen. The

present experiment demonstrated improved motility, membrane integrity, survival, and normality of spermatozoa and their acrosomes after cold storage for up to 72 h of roosters' sperms with garlic antioxidants that scavenge ROS in the sperm cell. Ultimately, the fertilizing ability of spermatozoa is most important and future studies will evaluate the effect of these antioxidants of garlic on this spermatozoa function.

References

- Al-Daraji, H.J. 1998. Effect of ascorbic acid supplementation on physiological and productive traits of Fawbro broiler breeder's flocks reared under hot climate. Ph. D. Dissertation, College of Agriculture, University of Baghdad.
- Al-Daraji, H.J. 2000. Effect of vitamin E on semen quality and fertilizing ability of roosters. *Dirasat, Agric. Sci.* **27**(3): 360 – 365.
- Al-Daraji, H.J. 2001. Effects of holding temperature and time on acrosomal abnormalities of fowl sperms. *Indian. J. Anim. Sci.* **71**(1): 32 – 34.
- Al-Daraji, H.J. 2002. Effect of vitamins A, C or E on quality of fowl semen stored for 24 hours at 4 °C. *Iraqi J. Agric.* **7**(6): 170-181.
- Al-Daraji, H.J. 2004. Diluent supplementation with vitamins A, C and E for improving fertilizing ability of indigenous roosters semen. Patent No. 3195, issued from C.O.S.Q.C., Iraq.
- Al-Daraji, H.J., Al-Tikriti, B.T.O., Hassan K.H. and Al-Rawi, A.A. 2002. A new techniques for determination of avian spermatozoa abnormalities. *Res. J. Bio. Tech.* **4**(1):47-64.
- Amagase, H., Schaffer, E.M. and Milner, J. 1996. Dietary components modify the ability of garlic to suppress 7, 12 – dimethyl (a) anthracene induced DNA adducts. *J. Nutr.* **126**: 817-824.
- Awazu, S. and Horie, T. 1997. Antioxidants in garlic. II. Protection of heart mitochondria by garlic extract and diallyl polysulfide from the doxorubicin – induced lipid peroxidation. Lanchance, P.P.ed. *Nutraceuticals Designer Foods III Garlic, Soy and Licorice* : 131-138 Food & Nutrition Press Trumbull, CT.
- Borek, C.2001. Antioxidant health effects of aged garlic extract. *J. Nutr.* **131**:1010S – 1015S.
- Borek, C.1997. Antioxidants and cancer. *Sci. Med.* **4**:51-62.
- Donoghue, A. and Donoghue, D.1997. Effects of water – and lipid – soluble antioxidants on turkey sperm viability, membrane integrity, and motility during liquid *Poultry Sci.* **76**: 1440-1445.
- El-Nasry, E., Khalil, H.M., Abaza M. and El-Saadany, A. 2004. Use of antioxidants in storing local cockerels semen 1. Effects on semen quality and fertility. WPC 2004, XXII World's Poultry Congress, Istanbul, Turkey.
- Geng, Z. and Lau, B.H.S.1997. Aged garlic extracts modulate glutathione redox cycle and superoxide dismutase activity in vascular endothelial cells. *Phytother. Res.* **11**: 54-56.
- Ide, N. and Lau, B.H.S.1997. Garlic compounds protect vascular endothelial cells from oxidized low density lipoprotein – induced injury. *J. Pharm. Pharmacol.* **49**:908-911.
- Imai, J., Ide, N., Nagae, S., Moriguchi, T., Matsuura, H. and Itakura, Y. 1994. Antioxidant and radical scavenging effects of aged garlic extract and its constituents. *Planta Med.* **60**:417-420.
- Kelso, K.A., Cerolini, S., Speake, B.K., Cavalchini, L.G. and Noble, R.C. 1997. Effects of dietary supplementation with α – linolenic acid on the phospholipid fatty acid composition and quality of spermatozoa in cockerel from 24 to 72

- weeks of age. *J. Reprod. Fert.* **110**: 53-59.
- Kelso, K.A., Cerolini, S., Noble, R.C., Sparks N.H.C. and Speake. B.K. 1996. Lipid and antioxidant changes in semen of broiler fowl from 25 to 60 weeks of age. *J.Reorod.Fert.***106**:201-206.
- Lake, P.E. and Stewart., J.M. 1978. Artificial insemination in poultry. Bulletin 213, Ministry of Agriculture, Fisheries and Food, London.
- Munday, J.S., James, A.K., Fray, L.M., Kirkwood, S.W and Thompson. K.G. 1999. Daily supplementation with aged garlic extract, but not raw garlic, protects low density lipoprotein against *in vitro* oxidation. *Atherosclerosis* **143**(2):399-404.
- Ohnishi, T. and Kojima. R. 1997. Antioxidant activities of aged garlic extracts and cancer chemotherapy. Lanchance, P.P. eds. *Nutraceuticals Designer Foods III Garlic, Soy and Licorice 1997*: 10 – 115 Food and Nutrition Press Trumbull, CT.
- Ohnishi, T. and Ohnishi. T. 2001. *In vitro* effects of aged garlic extract and other nutritional supplements on sickle erythrocytes. *J. Nutr.* **131**:1085S – 1092S.
- Pizzorri, J., Murray N.D. and Michael. T. 1999. *Textbook of Natural Medicine*, second ed. Churchill Livingstone, 1999.
- Reeve, V.E., Bosnic, M., Rosinova E. and Boehm-Wilcox. C. 1993. A garlic extract protects from ultraviolet B (280 – 320 nm) radiation induced suppression of contact hypersensitivity. *Photochem. Photobiol.* **58**:813-817.
- Ryu, K., Ide, N., Matsuura H. and Itakura. Y .2001. N⁶-91 – Deoxy – D – fructose – 1 – yl) – L – Arginine, an antioxidant compound identified in aged garlic extract. *J.Nutr.***131**: 972S – 976S.
- SAS.1996. SAS User's Guide: Statistics Version 6 th edn. SAS Inc., Cary, NC.
- Sexton, T.J.1976. Studies on the dilution of turkey semen. *Br. Poultry Sci.***17**: 179 – 186.
- Silagy, C.S. and Haw. N. 1994. Garlic as lipid lowering agent – a Meta analysis. *The Journal of the Royal College of Physicians.* **28**(1):39-45.
- Wei, Z. and Lau. B.H.S.1998. Garlic inhibits free radical generation and augments antioxidant enzyme activity in vascular endothelial cells. *Nutr.Res.***18**; 61 – 70.
- Wishart, G.J.1989. Physiological changes in fowl and turkey spermatozoa during *in vitro* storage. *Br. Poultry Sci.* **30**:443-454.
- Zaniboni, L., Parodi, L., Madjian, A., Gliozzi, T., Pizzi F. and Cerolini S..2004. Combined effect of DHA and α – tocopherol enrichment on quality and susceptibility to oxidation in chicken spermatozoa. WPC 2004, XXII World,s Poultry Congress, Istanbul, Turkey.