



Molecular Detection of Virulence Genes in *Aeromonas hydrophila* Isolated from Market Milk

Suneel Singh Gurjar¹, Abhishek Gaurav^{1*}, Deepak Kumar Sharma², Sudeep Solanki² and Vemula Nita Ida¹

¹Department of Veterinary Public Health and Epidemiology, College of Veterinary and Animal Science, Navania, Vallabhnagar, Udaipur, Rajasthan, INDIA

²Department of Veterinary Microbiology, College of Veterinary and Animal Science, Navania, Vallabhnagar, Udaipur, Rajasthan, INDIA

*Corresponding author: A Gaurav; E-mail: gaurav.vets@gmail.com

Received: 20 Feb., 2026

Revised: 24 March, 2026

Accepted: 29 March, 2026

ABSTRACT

Virulence gene profiling of *Aeromonas* species can be conducted using PCR to target various virulence factors, including endotoxins, enterotoxins, adhesins, cytotoxins, hemolysins, lipases, and proteases. A polymerase chain reaction (PCR) assay was employed for the detection of *Aeromonas hydrophila* (*A. hydrophila*) in raw market milk samples. A total of 350 milk samples were collected from different markets in Udaipur during the period from June 2025 to September 2025. PCR amplification using species-specific primers targeting the *gyrB* gene produced specific amplicons of 1100 bp, confirming the presence of *A. hydrophila*. The primers used in the study successfully amplified the corresponding target sequences of *gyrB* gene. Furthermore, several virulence-associated genes were amplified to assess the pathogenic potential of the isolates. The virulence genes targeted included *aer*, *ast*, *alt* and *act*. PCR amplification of these virulence genes yielded specific bands of 252 bp, 212 bp, 442 bp, and 232 bp, respectively. Out of the 12 *A. hydrophila* isolates, 11, 6, 9 and 8 isolates were found to be positive for *aer* (91.66%), *ast* (50%), *alt* (75%), and *act* (66.66%) genes, respectively. As raw milk can act as a vehicle for transmission of pathogenic strains to humans, its contamination represents a significant public health risk. The presence of virulence genes among the recovered isolates of *A. hydrophila* in this study highlights the pathogenic potential of milk-borne microorganisms, emphasizing the need for continuous surveillance through molecular approaches to strengthen food safety management and prevent future outbreaks.

HIGHLIGHTS

- ❶ The confirmation of *Aeromonas hydrophila* in raw market milk samples collected from Udaipur.
- ❷ Virulence gene profiling revealed a high prevalence of pathogenic determinants among isolates.
- ❸ The occurrence of virulent *A. hydrophila* in raw milk highlights the importance of continuous molecular surveillance for effective food safety management.

Keywords: *Aeromonas hydrophila*, PCR, virulence genes, milk, Udaipur

The genus *Aeromonas* has undergone several taxonomic and nomenclatural revisions over time. Initially classified within the family Vibrionaceae alongside the genera *Vibrio*, *Photobacterium*, and *Plesiomonas*, advances in molecular and phylogenetic analyses later resulted in its reclassification into the family Aeromonadaceae, which comprises mesophilic and psychrotrophic groups. *Aeromonas* species are widely distributed in aquatic environments and are increasingly recognized as emerging

foodborne and opportunistic pathogens affecting both humans and animals, particularly fish. In humans, these bacteria are most commonly associated with gastroenteritis, although they are also implicated in several extra-

How to cite this article: Gurjar, S.S., Gaurav, A., Sharma, D.K., Solanki, S. and Ida, V.N. (2026). Molecular Detection of Virulence Genes in *Aeromonas hydrophila* Isolated from Market Milk. *J. Anim. Res.*, **16**(02): 73-78.

Source of Support: None; **Conflict of Interest:** None



intestinal infections. While contaminated water remains the principal route of transmission, foodborne exposure has emerged as an important public health concern (Teodoro *et al.*, 2022). In India, *Aeromonas*-associated gastroenteritis is frequently reported, and studies from Kolkata have demonstrated the enterotoxic potential of *Aeromonas* species and their prevalence in food sources such as meat, milk, and fish (Manna *et al.*, 2013).

Several species of *Aeromonas* are considered significant human pathogens capable of causing a broad spectrum of diseases, including primary and secondary septicemia, particularly among immunocompromised individuals. They are also associated with severe wound infections, including those occurring during therapeutic leech therapy, as well as infections in otherwise healthy individuals. In addition, *Aeromonas* species have been linked to meningitis, peritonitis, respiratory tract infections, and infections involving bones, joints, and eyes (Janda and Abbott, 1998; Fernandez-Escartin and Garcia, 2001). These bacteria are believed to contribute significantly to the globally recognized seasonal illness known as “summer diarrhea”, which predominantly affects travelers, children under five years of age, and elderly individuals. The pathogenicity of *Aeromonas* species is attributed to the production of multiple virulence factors, including flagella, adhesins, elastase, hemolysins, aerolysins, phospholipases, enterotoxins, and lipases, which facilitate colonization, invasion, and tissue damage in the host (Carusi *et al.*, 2024).

Virulence profiling of *Aeromonas* species can be effectively performed using PCR-based detection of genes encoding toxins, adhesins, hemolysins, lipases, and proteases. Among these, exotoxins constitute the major virulence

determinants, including hemolysin (*hlyA*), aerolysin (*aerA*), cytotoxic heat-labile enterotoxin (*act*), cytotoxic heat-labile enterotoxin (*alt*), and cytotoxic heat-stable enterotoxin (*ast*) (Roges *et al.*, 2020). As Gram-negative bacteria with increasing clinical and environmental significance, *Aeromonas* species warrant continuous surveillance and effective control measures (Selim *et al.*, 2025). Moreover, the genus occupies a critical position within the One Health framework and may play an important role in the dissemination of antimicrobial resistance in the environment (Bojar *et al.*, 2025). Considering the growing public health importance of *A. hydrophila* in raw market milk, the present investigation was undertaken to detect virulence genes among *A. hydrophila* isolates.

MATERIALS AND METHODS

A total of 350 pooled raw milk samples were randomly collected from retail dairies across different areas of Udaipur city, Rajasthan from June 2025 to September 2025. The samples which were found to be positive for *A. hydrophila* based on cultural identification, Gram’s staining, motility characteristics and biochemical confirmation were subjected to molecular characterization based on the detection of virulence genes.

Detection of virulence genes in *A. hydrophila* isolates

Isolation of DNA from pure test culture was undertaken by using the Nucleo-pore gDNA fungal/bacterial mini kit (Genetix Biotech Asia Pvt. Ltd, India). The primers used for the molecular characterization of *A. hydrophila* isolates targeting the species-specific gene (*gyrB*) and virulence genes (*aer*, *ast*, *alt* and *act*) are described in Table 1.

Table 1: Primers used for molecular characterization of *A. hydrophila* isolates targeting species specific and virulence genes

Sl. No.	Target Gene	Sequence	Product Size (Bp)	Reference
1	<i>gyrB</i>	F: TCCGGCGGTCTGCACGGCGT R: TTGTCCGGGTGTACTCGTC	1100	Algammal <i>et al.</i> (2020) Xu <i>et al.</i> (2023)
2	<i>Alt</i>	F: TGACCCAGTCCTGG R: GGTGATCGATCACC	442	Hu <i>et al.</i> (2012) Mansour <i>et al.</i> (2019)
3	<i>ast</i>	F: CGCCATCAACAGCTCGCCCA R: CGGGCCTCGTTGAGGAAGCG	212	Bernabè <i>et al.</i> (2023) Sadiq <i>et al.</i> (2024)
4	<i>aer</i>	F: GCAGAACCCATCTATCCAG R: TTTCTCCGGTAACAGGATTG	252	Porteen <i>et al.</i> (2006) Narmada (2022)
5	<i>act</i>	F: AGAAGGTGACCACCACCAAGAACA R: AACTGACATCGGCCTGAACTC	232	Abo-Shama <i>et al.</i> (2025) Eid <i>et al.</i> (2019)

The PCR procedures to screen the species-specific and virulence genes were standardized. Followed by preliminary trials, the reaction mixture was optimized to contain 12.5µl Green Taq PCR master mix, 10 nmol (0.5µl) of each forward and reverse primer, 10.5µl nuclease free water and 1µl of DNA template. The PCR reaction was carried out in a thermal cycler with a pre-heated lid set to 105°C. The cycling conditions for the various primer pairs are detailed in (Table 2).

A 1% agarose gel was prepared in 1X TBE buffer, stained with ethidium bromide, and cast using an acrylic comb. The gel was run in 1X TBE buffer, with wells oriented toward the cathode. PCR products (6 µl) and a 100 bp/1 kb DNA ladder (5 µl) were loaded into the wells, while a negative control was included in the last well. Electrophoresis was performed at 120 V for 60 min, and amplified products were visualized under a UV transilluminator.

RESULTS AND DISCUSSION

All the 12 *A. hydrophila* isolates obtained from 350 raw milk samples were confirmed by using the primers targeting the species-specific (*gyrB*) gene, yielding the

product size of 1100 bp, (Fig. 1). Similarly, *gyrB* gene was used to confirm the *A. hydrophila* isolates in earlier studies conducted by Yanez *et al.* (2003), Soler *et al.* (2004) & Ninh *et al.* (2021). Soler *et al.* (2004) used *gyrB* & *rpo D* gene sequence independently as excellent molecular markers for assessing phylogeny in the genus *Aeromonas*.




Fig. 1: Agarose gel showing PCR amplified product (1100 bp) for *gyrB* gene. M=1kb DNA ladder, positive samples (L1= 1, L2=2, L3=3, L4=4, L5= 5, L6=6, NC= negative control)

PCR assay for the detection *alt*, *ast*, *aer* and *act* genes associated with virulence in *A. hydrophila* were standardized with primers for *alt* (Mansour *et al.*, 2019 & Hu *et al.*, 2012); *ast* (Sadiq *et al.*, 2024 & Bernabe *et al.*, 2023); *aer* (Narmada, 2022 & Porteen *et al.*, 2006) and

Table 2: Steps and conditions of thermal cycling for different primer pairs in PCR

Primers	Cycling conditions				
	Initial denaturation	Denaturation	Annealing	Extension	Final extension
<i>gyrB</i>	94°C	94°C	62°C	72°C	72°C
	5 min	30 sec	30 sec	1 min	5 min
Repeated for 30 cycles					
<i>alt</i>	95°C	94°C	55°C	72°C	72°C
	5 min	30 sec	40 sec	45 sec	5 min
Repeated for 30 cycles					
<i>ast</i>	95°C	94°C	60°C	72°C	72°C
	5 min	30 sec	30 sec	1 min	5 min
Repeated for 30 cycles					
<i>aer</i>	94°C	94°C	55°C	72°C	72°C
	5 min	30 sec	50 sec	1 min	5 min
Repeated for 30 cycles					
<i>act</i>	94 °C	94 °C	53°C	72°C	72°C
	5 min	30 sec	40 sec	30 sec	5 min
Repeated for 30 cycles					

act (Abo-Shama et al., 2025 & Eid et al., 2019) genes. The PCR assay showed specific amplification products of 442bp, 212bp, 252 bp and 232 bp for *alt*, *ast*, *aer* and *act* genes, respectively (Fig. 2 to Fig. 5).



Fig. 2: Agarose gel showing PCR amplified product (442 bp) for *alt* gene. M=100bp DNA ladder, positive samples (L1= 1, L2=2, L3=4, L4=6, L5=7, L6=9, NC= negative control)




Fig. 3: Agarose gel showing PCR amplified product (212 bp) for *ast* gene. M=100bp DNA ladder, positive samples (L1= 1, L2=3, L3=4, L4=6, L5=7, L6= 11, NC= negative control)




Fig. 4: Agarose gel showing PCR amplified product (252 bp) for *aer* gene. M=100 bp DNA ladder, positive samples (L1= 1, L2=2, L3=3, L4=4, L5=5, L6=6, NC= negative control)




Fig. 5: Agarose gel showing PCR amplified product (232 bp) for *act* gene. M=100bp DNA ladder, positive samples (L1= 1, L2= 2, L3=3, L4= 5, L5=6, L6=8, NC= negative control)

In the current study, out of the 12 *A. hydrophila* isolates 9, 6, 11 and 8 isolates were found to be positive for *alt* (75%), *ast* (50%), *aer* (91.66%) and *act* (66.66%) genes, respectively. The PCR amplified products of *A. hydrophila* isolates revealed band size of 442 bp for *alt*, 212 bp for *ast*, 252 bp for *aer*, and 232 bp for *act* gene by agarose gel electrophoresis. Various virulence genes in *A. hydrophila* were targeted in the present study viz. *aer*, *alt*, *act* & *ast* revealing prevalence as 91.66%, 75%, 66.66% & 50%, respectively. Slightly lower prevalence of *aer* gene was reported by Sughra et al. (2022) & Yano et al. (2015) as 75% & 74%, respectively. While, lower prevalence rate of *aer* gene in *A. hydrophila* were reported by El-Baz and Hendam (2022), Ayoub et al. (2024) & Saleh et al. (2021) as 42.1%, 57.9% & 40%, respectively. Yano et al. (2015), Saleh et al. (2021) & Ayoub et al. (2024) reported lower prevalence rate of *alt* gene as 30%, 30% & 26.3%, respectively. While, 100% prevalence of *alt* gene was reported by Rather et al. (2014) among *A. hydrophila* isolates.

Pathogenicity in *A. hydrophila* is multifactorial and largely mediated by toxins such as *ast*, *alt*, aerolysin-hemolysin, and the *act*-encoded cytotoxic enterotoxin, which are associated with human gastroenteritis (Balsalobre et al., 2009). The virulence of *A. hydrophila* is closely linked to the production of several virulence factors and its biofilm-forming ability, including serine protease, hemolysin, aerolysin, cytotoxic enterotoxin, and adhesins such as fimbriae and S-layer proteins (Zhou et al., 2019). Exotoxins, including the cytotoxic heat-labile enterotoxin (*act*), cytotoxic heat-labile enterotoxin (*alt*), cytotoxic heat-stable enterotoxin (*ast*), aerolysin/hemolysin, lipase, and phospholipase, play important roles in pathogenicity (Kingombe et al., 2010). Although several PCR methods have been developed for detecting individual cytotoxin and hemolysin genes, a multiplex PCR assay was designed to simultaneously amplify the hemolysin and aerolysin genes of *A. hydrophila* and *A. sobria* (Wang et al., 2003). Variations in the frequency of virulence genes among *A. hydrophila* isolates indicate diversity in their virulence gene profiles.

ACKNOWLEDGMENTS

The authors are thankful to the Dean, College of Veterinary and Animal Science, Navania, Vallabh Nagar, Udaipur for providing all the facilities during the research period.

REFERENCES

- Abo-Shama, U.H., El Raheem, A.A.A., Alsaadawy, R.M. and Sayed, H.H. 2025. Phenotypic and genotypic characterization of *Aeromonas hydrophila* isolated from freshwater fishes at Middle Upper Egypt. *Sci. Rep.*, **15**(1): 01-11.
- Algammal, A.M., Mohamed, M.F., Tawfik, B.A., Hozzein, W.N., El Kazzaz, W.M. and Mabrok, M. 2020. Molecular typing, antibiogram, and PCR-RFLP-based detection of *Aeromonas hydrophila* complex isolated from *Oreochromis niloticus*. *Pathogens*, **9**(3): 238-250.
- Ayoub, H.F., Khafagy, A.R., Esawy, A.M., El-Moaty, N.A., Alwutayd, K.M., Mansour, A.T. and El-Tarabili, R.M. 2024. Phenotypic, molecular detection, and Antibiotic Resistance Profile (MDR and XDR) of *Aeromonas hydrophila* isolated from Farmed Tilapia zillii and Mugil cephalus. *BMC Vet. Res.*, **20**(1): 01-13.
- Balsalobre, L.C., Dropa, M., Matte, G.R. and Matte, M.H. 2009. Molecular detection of enterotoxins in environmental strains of *Aeromonas hydrophila* and *Aeromonas jandaei*. *J. Water Health*, **7**(4): 685-691.
- Bernabè, G., Brun, P., Pietra, G.D., Zatta, V., Asad, S., Meneghello, S. and Castagliuolo, I. 2023. Prevalence and virulence potential of *Aeromonas* spp. isolated from human diarrheal samples in North East Italy. *Microbiol. Spectr.*, **11**(6): 1-20.
- Bojar, B.E., Craig III, A.T., Leduc, A., Blumenthal, M., Mayo, B., Ahmed, A.S., Cahak, C., Beattie, R. and Skwor, T. 2025. Similar antimicrobial resistance and virulence profiles among *Aeromonas* isolates from recreational beaches, post-chlorinated wastewater and clinical samples in Milwaukee, Wisconsin USA. *Sci. Total Environ.*, **970**: 179055.
- Carusi, J., Kabuki, D.Y., De Seixas Pereira, P.M. and Cabral, L. 2024. *Aeromonas* spp. in drinking water and food. Occurrence, virulence potential and antimicrobial resistance. *Food Res. Int.*, **175**: 113710.
- Eid, H., Soliman, Z. and Hanafy, A.S. 2019. Molecular detection of some virulence genes in *Aeromonas* species isolated from fishes and water of Manzala Lake. *Suez Canal Vet. Med. J.*, **24**(2): 231-243.
- El-Baz, A. and Hendam, B. 2022. Virulence genes and antibiotic resistance of *Aeromonas hydrophila* isolated from marketed milk. *Veterinarija ir Zootechnika*, **80**(1): 10-19.
- Fernandez-Escartin, E. and Garcia, S. 2001. Guide to Foodborne Pathogens. In: Miscellaneous agents *Brucella*, *Aeromonas*, *Plesiomonas* and b-haemolytic *streptococci*, John Wiley and Sons, New York, USA, pp 295-313.
- Hu, M., Wang, N., Pan, Z.H., Lu, C.P. and Liu, Y.J. 2012. Identity and virulence properties of *Aeromonas* isolates from diseased fish, healthy controls and water environment in China. *Lett. Appl. Microbiol.*, **55**(3): 224-233.
- Janda, J.M. and Abbott, S.L. 1998. Evolving concepts regarding the genus *Aeromonas*: An expanding panorama of species, disease presentations, and unanswered questions. *Clin. Infect. Dis.*, **27**(2): 332-344.
- Kingombe, C.I.B., Huys, G., Tonolla, M., Albert, M.J., Swings, J., Peduzzi, R. and Jemmi, T. 1999. PCR detection, characterization, and distribution of virulence genes in *Aeromonas* spp. *Appl. Environ. Microbiol.*, **65**(12): 5293-5302.
- Manna, S.K., Maurye, P., Dutta, C. and Samanta, G. 2013. Occurrence and virulence characteristics of *Aeromonas* species in meat, milk and fish in India. *J. Food Saf.*, **33**(4): 461-469.
- Mansour, A., Mahfouz, N.B., Husien, M.M. and El-Magd, M.A. 2019. Molecular identification of *Aeromonas hydrophila* strains recovered from Kafrelsheikh fish farms. *Slov. Vet. Res.*, **56**(22): 201-208.
- Narmada, S. 2022. Molecular characterization, virulence gene profiling and multidrug resistance profiles of *Aeromonas hydrophila* isolated from raw fish samples collected in and around Tirupati, Andhra Pradesh, India. *J. Pharm. Innov.*, **11**(5): 1669-1673.
- Nhinh, D.T., Le, D.V., Van, K.V., Huong Giang, N.T., Dang, L.T. and Hoai, T.D. 2021. Prevalence, virulence gene distribution and alarming the multidrug resistance of *Aeromonashydrophila* associated with disease outbreaks in freshwater aquaculture. *ATB.*, **10**(5): 01-17.
- Porteen, K., Agarwal, R.K. and Bhilegaonkar, K.N. 2006. PCR-based detection of *Aeromonas* from milk samples. *J. Food Technol.*, **4**(2): 111-115.
- Rather, M.A., Willayat, M.M., Wani, S.A., Munshi, Z.H. and Hussain, S.A. 2014. A multiplex PCR for detection of enterotoxin genes in *Aeromonas* species isolated from foods of animal origin and human diarrhoeal samples. *J. Appl. Microbiol.*, **117**(6): 1721-1729.
- Roges, E.M., Goncalves, V.D., Cardoso, M.D., Festivo, M.L., Siciliano, S., Berto, L.H., Pareria, V.L.D.A., Rodrigues, D.D.P. and Aquino, M.H.C.D. 2020. Virulence associated genes and antimicrobial resistance of *Aeromonas hydrophila* isolates from animal, food and human sources in Brazil. *Biomed Res. Int.*, **8**(1): 01-08.
- Sadiq, S.T., Al-Hamdani, A.H.A. and Taha, Z.M.A. 2024. Clonal dispersion and pathogenic potential of multidrug-resistant *Aeromonas* spp. isolated from *Oncorhynchus mykiss* with haemorrhagic septicaemia. *Vet. Res. Forum.*, **15**(10): 529-536.
- Saleh, A., Elkenany, R. and Younis, G. 2021. Virulent and multiple antimicrobial resistance *Aeromonas hydrophila* isolated from diseased Nile tilapia fish (*Oreochromis*



- niloticus*) in Egypt with sequencing of some virulence-associated genes. *Biocontrol Sci.*, **26**(3): 167-176.
- Selim, S., Harun-Ur-Rashid, M., Jahan, I. and Elamir, M.Y.M. 2025. Antibiotic susceptibility patterns of *Aeromonas*: Insights from aquatic ecosystems to public health impacts. *Egypt. J. Aquat. Res.*, **51**(2): 257-269.
- Soler, L., Yanez, M.A., Chacon, M.R., Aguilera-Arreola, M.G., Catalan, V., Figueras, M.J. and Martinez-Murcia, A.J. 2004. Phylogenetic analysis of the genus *Aeromonas* based on two housekeeping genes. *Int. J. Syst. Evol. Microbiol.*, **54**(5): 1511-1519.
- Sughra, F., Hafeez-ur-Rehman, M., Abbas, F., Altaf, I., Hassan, Z., Bhatti, A. and Ali, K. 2022. Molecular characterisation and genetic analysis of aerolysin and haemolysin in *Aeromonashydrophila* isolated from diseased Labeorohita by polymerase chain reaction. *J. Fish.*, **10**(3): 103402-103402.
- Teodoro, J.R., Carvalho, G.G., Queiroz, M.M., Levy, C.E. and Kabuki, D.Y. 2022. Incidence, evaluation of detection and identification methods and antimicrobial resistance of *Aeromonas* spp. in ready-to-eat foods. *Int. J. Food Microbiol.*, **379**:109862.
- Wang, G., Clark, C.G., Liu, C., Pucknell, C., Munro, C.K., Kruk, T.M. and Rodgers, F.G. 2003. Detection and characterization of the hemolysin genes in *Aeromonashydrophila* and *Aeromonassobria* by multiplex PCR. *J. Clin. Microbiol.*, **41**(3): 1048-1054.
- Xu, T., Rasmussen-Ivey, C.R., Moen, F.S., Fernandez-Bravo, A., Lamy, B., Beaz-Hidalgo, R. and Liles, M.R. 2023. A global survey of hypervirulent *Aeromonas hydrophila* (vAh) identified vAh strains in the lower Mekong river basin and diverse opportunistic pathogens from farmed fish and other environmental sources. *Microbiol. Spectr.*, **11**(2): 01-15.
- Yanez, M.A., Catalan, V., Apraiz, D., Figueras, M.J. and Martinez-Murcia, A.J. 2003. Phylogenetic analysis of members of the genus *Aeromonas* based on *gyrB* gene sequences. *Int J Syst Evol Microbiol.*, **53**(3): 875-883.
- Yano, Y., Hamano, K., Tsutsui, I., Aue-Umneoy, D., Ban, M. and Satomi, M. 2015. Occurrence, molecular characterization, and antimicrobial susceptibility of *Aeromonas* spp. in marine species of shrimps cultured at inland low salinity ponds. *Food Microbiol.*, **47**: 21-27.
- Zhou, Y., Yu, L., Nan, Z., Zhang, P., Kan, B., Yan, D. and Su, J. 2019. Taxonomy, virulence genes and antimicrobial resistance of *Aeromonas* isolated from extra-intestinal and intestinal infections. *BMC Infect. Dis.*, **19**(1): 01-09.